رسیلاست®۱۰
بیلاستین

قرص بازشونده در دهان بیلاستین ۱۰
بیلاستین به شکل قرص‌های بازشونده در دهان حاوی ۱۰ میلی‌گرم بیلاستین ارائه می‌شوند. ۱۰ قرص در هر بلیستر و ۳ بلیستر در یک جعبه بسته بندی می شود.

دسته دارویی: آنتی هیستامین (گیرنده H_1)

هشدارها و احتیاطات:
- در صورت سابقه حساسیت به بیلاستین یا سایر آنتی هیستامین ها (مانند واکنش های آنافیلاکتیک) باید مصرف بیلاستین قطع و به پزشک یا داروساز اطلاع داده شود.
- در صورت نارسایی کلیوی متوسط تا شدید ، پیش از مصرف این دارو با پزشک یا داروساز مشورت کنید.
- هنگام مصرف این دارو، از رانندگی و فعالیت های نیازمند هوشیاری کامل، خودداری نمایید.
- دارو باید دقیقا طبق دستور پزشک یا داروساز مصرف شود.
- بیلاستین نباید در بیماران با سابقه QT prolongation مانند افراد با سابقه آریتمی قلبی، هایپوکالمی؛ هایپومنیزیمی؛ برادی کاردی شدید؛ سابقه خانوادگی ایست ناگهانی قلبی؛ همزمان با سایر داروهای تشدیدکننده QT prolongation، استفاده شود.
- مصرف دارو در کودکان کمتر از ۶ سال و همچنین با وزن کمتر از ۲۰ کیلوگرم ممنوع است.

مقدار و نحوه مصرف:
- دارو را یک ساعت قبل یا دو ساعت بعد از غذا با آب میوه مصرف کنید.
- اگر دوز روزانه فراموش شد، به محض یادآوری مصرف شود و دوز بعدی در زمان معمول طبق دستور پزشک مصرف شود. برای جبران دوز فراموش شده نباید دوبرابر مصرف شود.
- در صورت اختلال در عملکرد کلیوی یا کبدی با پزشک مشورت شود.
- قرص بازشونده در دهان را در دهان قرارداده تا در بزاق دهان به سرعت باز شده و به راحتی بلعیده شود. همچنین می توان قرص را در یک قاشق آب باز کرد. اطمینان حاصل کنید که پس از مصرف، باقی مانده ای از قرص در قاشق نمانده باشد. دقت شود برای باز کردن قرص از هیچ آب میوه ای بجای آب استفاده نشود.

مقدار مصرف هر دارو توسط پزشک تعیین می گردد اما دستور مصرف معمول این دارو به شرح زیر است:
اختلالات آلرژیک (التهاب ملتحمه بینی و کهیر) :
- کودکان ۶ تا ۱۱ ساله (وزن بدن حداقل ۲۰ کیلوگرم):
۱۰ میلی گرم بیلاستین (۱ قرص بازشونده در دهان) یک بار در روز

عوارض جانبی:
هر دارو به موازات اثرات درمانی ممکن است باعث بروز بعضی عوارض ناخواسته نیز شود؛ اگر چه همه این عوارض در یک فرد دیده نمی شود؛ در صورت بروز عوارض زیر و یا هرگونه عارضه ی دیگر، با پزشک یا داروساز مشورت نمایید:
عوارض نیازمند مراقبت پزشکی: در صورت بروز واکنش های آلرژیک مانند مشکل در تنفس، سرگیجه، افتادن یا از دست دادن هوشیاری، تورم صورت، لب ها، زبان یا گلو و یا تورم و قرمزی پوست، درمان را متوقف و با پزشک تماس گرفته شود.
عوارض رایج: سردرد، سرگیجه
عوارض غیر رایج: نوار قلب غیر طبیعی، ضربان قلب نامنظم؛ تغییرات در عملکرد کبد و کلیه؛ اختلال در خواب، اضطراب، خستگی، سرگیجه؛ دل درد، افزایش اشتها، افزایش وزن، حالت تهوع، اسهال، التهاب معده، سو، هاضمه، تشنگی، احساس ضعف؛ تنگی نفس، خشکی دهان، بینی خشک یا ناراحت کننده؛ خارش؛ تبخال در دهان؛ وزوز گوش ؛ افزایش چربی خون

تداخلات دارویی:
اگر داروی دیگری مصرف می کنید یا ممکن است مصرف کنید، پزشک یا داروساز خود را مطلع نمایید.
داروهایی که ممکن است با بیلاستین تداخل داشته باشند:
سیکلوسپورین؛ اریترومایسین؛ کتوکونازول؛ ریتونوایر؛ ریفامپیسین؛ دیلتیازم؛ لورازپام

شرایط نگهداری:
دارو را در دمای زیر ۳۰ درجه سانتیگراد و دور از نور و رطوبت نگهداری نمایید.

ساخت شرکت داروسازی پور سینا
تهران – ایران
PURSINA
پورسینا

Resbilast® 10
Bilastine

Bilastine orally disintegrating tablet (ODT) are presented as forms containing 10 mg Bilastine; 10 tablets per blister pack, 3 blister with a leaflet in a box.

Category: Antihistamine (H1-Receptor)

Precautions and warnings:

- If there is allergic history (such as anaphylactic reactions) to Bilastine or other antihistamines, stop taking Bilastine.
- Talk to doctor or pharmacist before using this tablet if there is moderate or severe renal impairment.
- Precaution is necessary before driving or operating machinery.
- Always use this medicine exactly as doctor or pharmacist has told.
- Bilastine should not be used in patients with a history of QTc prolongation like patients with history of cardiac arrhythmias; hypokalemia; hypomagnesemia; significant bradycardia; family history of sudden cardiac death; concomitant use of other QT/QTc-prolonging drugs.
- Do not give Bilastine to children under 6 years old with a body weight below 20 kg.

Dosage and administration:

- Take the tablet one hour before or two hours after taking food or fruit juice.
- If you forget to take the daily dose on time, give it on the same day as soon as you remember. Then, give the next dose on the next day at the usual time as prescribed by the doctor. Do not take a double dose to make up for a forgotten dose.
- If you are suffering from kidney or liver disorders, consult the doctor.
- The orally disintegrating tablet is to be placed in the mouth where it disperses rapidly in saliva, so it can be easily swallowed.
- Alternatively, the orodispersible tablet may be dispersed in a teaspoon of water before administration. It has to be ensured that no rest of sediment remains in the spoon. fruit juices should not be used for dispersion.

Dosage of each drug is determined by doctor, but the usual dosage of Bilastine is as follows:
Allergic disorders (rhinoconjunctivitis and urticaria):

- Children 6 to 11 years old (body weight of at least 20 kg):

10 mg Bilastine (1 orodispersible tablet) once a day

Adverse reactions:

Like all medicines, this medicine may cause unwanted side effects. In case of the following side effects, consult with the

pharmacist or doctor:

Those need for medical attention:
Stop treatment and contact doctor if you have allergic reactions like difficulty in breathing, dizziness, collapsing or losing consciousness, swelling of your face, lips, tongue or throat, and/or swelling and redness of the skin

- Common: Headache; drowsiness
- Uncommon: Abnormal ECG heart tracing, irregular heartbeat; changes in liver or kidney function; dizziness, anxiety, tiredness, feeling of weakness, difficulty in sleeping; stomach pain, increased appetite, increased weight, nausea, indigestion, diarrhea, gastritis, thirst; dyspnea; dry or uncomfortable nose, dry mouth; itching; oral herpes; fever; tinnitus; increased blood fat

Drug interactions:

- Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.
- Medicines which may interact with Bilastine:

Cyclosporine; Erythromycin; Ketoconazole; Ritonavir; Rifampicin; Diltiazem; Lorazepam

Storage:

Store below 30° C, protect from light and moisture.

Manufactured by
Pursina Pharmaceutical Co.
Tehran-Iran
PURSINA
پورسینا